

## 論文

- Hori, K., L-Y, Kuo., W-L, Chiou., A, Ebihara. and N. Murakami (2016). The geographical distribution of sexual and apogamous types of *Dryopteris formosana* and *Dryopteris varia* (Dryopteridaceae) in Taiwan. Acta Phytotax. Geobot 68: 23-32.
- Hori, K., Watano, Y. and N. Murakami (2016). Hybrid origin of the apogamous fern species *Dryopteris hondoensis* (Dryopteridaceae). Acta Phytotax. Geobot 67: 133-146.
- Hori, K., Matsumoto, M., Nakato, N., Ebihara, A. and N. Murakami (2015). Geographical distribution of sexual and apogamous types of *Dryopteris chinensis* (Dryopteridaceae) in Japan. Acta Phytotax. Geobot. 66: 35-45.
- Hori, K., Ebihara, A., Nakato, N. and N. Murakami (2015). *Dryopteris protobissetiana* (Dryopteridaceae), a new diploid sexual species of the *Dryopteris varia* complex (subg. *Erythrovariae*, sect. *Variae*) from Yakushima Island, Kagoshima, Japan. Acta Phytotax. Geobot. 66: 47-57.
- Hori, K., Tono, A., Fujimoto, K., Kato, J., Ebihara, A., Watano, Y. and N. Murakami (2014). Reticulate evolution in the apogamous *Dryopteris varia* complex (Dryopteridaceae, subg. *Erythrovariae*, sect. *Variae*) and its related sexual species in Japan. J. Plant Res. 127: 661-684.

## 他1編

## 学会発表

- Hori, K., Okuyama, Y., Watano, Y., & N. Murakami (2016) Reticulate evolution maintaining original genome constitutions in the *Dryopteris varia* complex. East Asian Plant Diversity and Conservation 2016. (口頭), 2016/8/23 (招待講演)

## 他17編